



RECOMMENDING MAXIGESIC TO YOUR DENTAL PATIENTS

IS/DOES YOUR PATIENT:

	YES	
▶ Under 12 years of age?	<input type="checkbox"/>	IF YES, DO NOT USE
▶ Pregnant or trying to become pregnant?	<input type="checkbox"/>	IF YES, DO NOT USE
▶ Breastfeeding or intending to breastfeed?	<input type="checkbox"/>	IF YES, DO NOT USE
▶ Have a stomach or duodenal ulcer?	<input type="checkbox"/>	IF YES, DO NOT USE
▶ Have impaired kidney or liver function?	<input type="checkbox"/>	IF YES, DO NOT USE
▶ Suffer from heart failure?	<input type="checkbox"/>	IF YES, DO NOT USE
▶ Allergic to paracetamol, ibuprofen, aspirin or other anti-inflammatory medicines?	<input type="checkbox"/>	IF YES, DO NOT USE
▶ Regularly drink large amounts of alcohol?	<input type="checkbox"/>	IF YES, DO NOT USE
▶ Have asthma?	<input type="checkbox"/>	USE ONLY ON DOCTOR'S ADVICE
▶ Aged 65 years or older?	<input type="checkbox"/>	USE ONLY ON DOCTOR'S ADVICE

If the patient answers **NO** to all of these questions, then MAXIGESIC is suitable to recommend.

Before recommending please review the full Product Information available at www.maxigesic.com.au/faqs.html

MAXIGESIC is indicated for the temporary relief of pain associated with:

- | | | |
|---------------------|---------------------|---|
| ▶ Headache | ▶ Dental procedures | ▶ Period pain |
| ▶ Migraine headache | ▶ Backache | ▶ Muscular pain |
| ▶ Tension headache | ▶ Sore throat | ▶ Rheumatic pain |
| ▶ Sinus pain | ▶ Arthritis | ▶ Aches & pains associated with colds and flu |
| ▶ Toothache | ▶ Tennis elbow | ▶ Also reduces fever |

DIRECTIONS FOR USE (adults & children over 12 years):

Take one to two tablets with a large glass of water.

Repeat dose every six hours, as required, up to a maximum of 8 tablets in 24 hours.

MAXIGESIC can also be dissolved in a glass of water, if required.

Do NOT take MAXIGESIC with other products containing paracetamol, ibuprofen, aspirin or other anti-inflammatory medicines.

Adults: Do not take MAXIGESIC for more than a few days at a time unless advised to by a doctor.

Children and adolescents 12 years+: Do not take MAXIGESIC for more than 48 hours at a time unless advised to by a doctor.

AFT Pharmaceuticals Pty Ltd, Sydney
ABN 29105636413
WEBSITE www.aftpharm.com

FREE PHONE 1800 AFT PHARM (238 742)
FREE FAX 1800 041 026
EMAIL receptionau@aftpharm.com

AFT pharmaceuticals
Working to improve your health